

Consent form: Comirnaty (Pfizer) COVID-19 vaccination



Before completing this form we recommend you read the information sheet on the COMIRNATY (Pfizer) COVID-19 Vaccine on the Department of Health website (www.health.gov.au).

Patient Information

People who have a COVID-19 vaccination have a much lower chance of getting sick from COVID-19.

The COVID-19 vaccination is free and you can choose whether to have the vaccination or not.

There are two brands of vaccine in use in Australia. Both are effective and safe. Comirnaty (Pfizer) vaccine is preferred over AstraZeneca COVID-19 vaccine for adults under 60 years of age.

You need to have two doses of the same brand of vaccine. The person giving you your vaccination will tell you when you need to have the second vaccination.

Medical experts have studied COVID-19 vaccines to make sure they are safe. Most side effects are mild. They may start on the day of vaccination and last for around 1-2 days. The more common side effects may include tenderness or pain at the injection site, headache, fever, fatigue, malaise, joint and muscle pain. As with any vaccine or medicine, there may be rare and/or unknown side effects.

A very rare side effect of blood clotting (thrombosis) with low platelet levels (thrombocytopenia) has been reported following vaccination with AstraZeneca. This is not seen after the Comirnaty (Pfizer) vaccine.

Cases of Myocarditis (inflammation of the heart muscle) and/or pericarditis (inflammation of the lining of the heart) have been reported as very rare side effects after mRNA COVID-19 vaccines (including Comirnaty (Pfizer)). Cases primarily occurred within 14 days after vaccination, more often in males aged under 30 years and after the second dose. COVID-19 Vaccine AstraZeneca is not associated with an increased risk of myocarditis/pericarditis.

Tell your healthcare provider if you have any side effects like a sore arm, chest pain, heart palpitations, shortness of breath, headache, fever, body aches or any symptom that is unusual for you. You may also be contacted after receiving the vaccine to see how you are feeling after your vaccination.

You will be asked to remain in the centre for 15 minutes observation after your vaccination, but if you have a previous history of anaphylaxis to a vaccine you will be required to wait for 30 minutes.

Some people may still get COVID-19 after vaccination, so you must still follow public health precautions as required in your State or Territory to stop the spread of COVID-19 including: physical distancing, wearing a mask, enhanced personal hygiene (hand washing/sanitising), staying at home if unwell with cold-like symptoms and promptly getting tested for COVID-19.

Vaccination providers record all vaccinations on the Australian Immunisation Register, as required by Australian Law. You can view this online in your Medicare/ MyGov/MyHealthRecord account.

For information on how your personal details are collected, stored and used visit <https://www.health.gov.au/covid19-vaccines>

On the day you receive your vaccine

Before you get vaccinated, tell the person giving you the vaccination if you:

- have had an allergic reaction, particularly anaphylaxis (a severe allergic reaction) to a previous dose of a COVID-19 vaccine, to an ingredient of a COVID-19 vaccine, or to any other vaccines or medications.
- are immune-compromised. This means that you have a weakened immune system that may make it harder for you to fight infections and other diseases.
- if you have a history of inflammatory cardiac illness (eg. pericarditis, myocarditis, endocarditis), acute rheumatic fever, dilated cardiomyopathy (for people under 30 years of age), congenital heart disease, severe heart failure or if you are a heart transplant recipient.

Pre-Vaccination Screening

Yes No

<input type="checkbox"/>	<input type="checkbox"/>	Do you have any serious allergies, particularly anaphylaxis, to anything and specifically a COVID vaccine ingredient?
<input type="checkbox"/>	<input type="checkbox"/>	Have you had anaphylaxis to another vaccine or medication?
<input type="checkbox"/>	<input type="checkbox"/>	Have you had a serious adverse event, that following expert review was attributed to a previous dose of a COVID-19 vaccine?
<input type="checkbox"/>	<input type="checkbox"/>	Have you ever had mastocytosis which has caused recurrent anaphylaxis?
<input type="checkbox"/>	<input type="checkbox"/>	Have you had COVID-19 before?
<input type="checkbox"/>	<input type="checkbox"/>	Do you have a bleeding disorder?
<input type="checkbox"/>	<input type="checkbox"/>	Do you take any medicine to thin your blood (an anticoagulant therapy)? If so, which medication do you take? _____
<input type="checkbox"/>	<input type="checkbox"/>	Do you have a weakened immune system (immunocompromised)?
<input type="checkbox"/>	<input type="checkbox"/>	Are you pregnant?
<input type="checkbox"/>	<input type="checkbox"/>	Have you been sick with a cough, sore throat, fever or are feeling sick in another way?
<input type="checkbox"/>	<input type="checkbox"/>	Have you had a COVID-19 vaccination elsewhere? If so, when: _____ and which one? <input type="checkbox"/> Astra Zeneca <input type="checkbox"/> Pfizer <input type="checkbox"/> Other, please specify: _____
<input type="checkbox"/>	<input type="checkbox"/>	Have you received any other vaccination in the last 7 days?
<input type="checkbox"/>	<input type="checkbox"/>	Have you ever fainted after a vaccination or are you especially scared of needles?
<input type="checkbox"/>	<input type="checkbox"/>	Have you ever had myocarditis or pericarditis?
<input type="checkbox"/>	<input type="checkbox"/>	Do you currently have, or have you recently had acute rheumatic fever or endocarditis?
<input type="checkbox"/>	<input type="checkbox"/>	Do you have congenital heart disease?
<input type="checkbox"/>	<input type="checkbox"/>	For people under 30 years of age: do you have dilated cardiomyopathy?
<input type="checkbox"/>	<input type="checkbox"/>	Do you have severe heart failure?
<input type="checkbox"/>	<input type="checkbox"/>	Are you a recipient of a heart transplant?

Please talk to your doctor if you have any questions or concerns before getting your COVID-19 vaccination.

Consent to receive COVID-19 vaccine

- I confirm I have received and understood information provided to me on Comirnaty (Pfizer) Covid-19 Vaccination
- I confirm that none of the conditions above apply, or I have discussed these and/or any other special circumstances with my regular healthcare provider and/or vaccination service provider
- I agree to receive a Comirnaty (Pfizer) Covid-19 Vaccine today

Patient's name: _____ Date of Birth: _____

Patient's signature: _____ Date: _____

- I am the patient's legal guardian or substitute decision maker, and agree to COVID-19 vaccination of the patient named above

Guardian/substitute decision maker's name: _____

Guardian/substitute decision maker's signature: _____ Date: _____

Doctor Use Only

- Nurse to administer COMIRNATY (Pfizer) 0.3mL (following dilution) IM

Doctor's Name: _____

Doctor's signature: _____ Date: _____